

## Folkestone Directional WaveRider Buoy

### Location

OS: 619711E 132538N  
 WGS84: Latitude: 51°03.5335'N Longitude: 01°08.2988'E

### Water Depth

12.7m CD

### Instrument Type

Datawell Directional WaveRider Buoy Mk III

### Data Quality

C1(%)	Sample interval
99	30 minutes

### Monthly Means

All times GMT

Month	H <sub>s</sub>	T <sub>p</sub>	T <sub>z</sub>	Direction	SST	No. of days
	(m)	(s)	(s)	(°)	(°C)	
January	0.89	5.9	3.8	168	9.4	31
February	0.69	6.0	3.7	158	8.4	28
March	0.59	6.5	3.9	149	9.2	31
April	0.33	5.4	3.4	118	10.4	30
May	0.59	5.2	3.6	148	12.9	31
June	0.49	4.8	3.4	150	15.3	30
July	0.62	4.5	3.4	175	16.8	30
August	0.44	4.7	3.5	145	18.1	31
September	0.46	5.3	3.8	142	17.6	30
October	0.45	5.1	3.5	132	15.3	31
November	0.61	5.6	4.0	154	11.6	30
December	0.77	5.7	3.7	152	9.2	31

Tables and plots of these values, together with the minimum and maximum values and the standard deviation are available on the website

Highest storm events in 2007									
Date/Time	H <sub>s</sub>	T <sub>p</sub>	T <sub>z</sub>	Dir.	Water level elevation (OD)	Tidal stage (hours re. HW)	Tidal range (m)	Tidal surge* (m)	Max. surge* (m)
08-Dec-2007 17:00	2.86	7.7	5.4	183	-2.2	HW - 6	4.5	-0.02	0.55
18-Jan-2007 13:00	2.59	-	4.4	-	-	HW + 3	4.9	-	-
06-Mar-2007 06:00	2.51	6.7	4.9	174	-1.87	HW + 5	5.6	0.16	0.81

\* Tidal information is obtained from the nearest recording tide gauge (the National Network gauge at Dover). The surge shown is the residual at the time of the highest H<sub>s</sub>. The maximum tidal surge is the largest positive surge during the storm event.

Year	Annual $H_s$ exceedance* (m)						Annual Maximum $H_s$	
	0.05%	0.5%	1%	2%	5%	10%	Date	$A_{max}$ (m)
2003	2.79	2.23	2.03	1.75	1.37	1.16	29-Nov-2003 13:30	3.07
2004	2.91	2.30	1.97	1.75	1.44	1.18	08-Oct-2004 12:00	3.25
2005	2.90	2.15	1.81	1.54	1.25	0.97	30-Dec-2005 14:00	3.15
2006	2.55	2.08	1.84	1.68	1.42	1.17	03-Dec-2006 09:00	3.13
2007	2.56	2.06	1.83	1.59	1.34	1.11	08-Dec-2007 17:00	2.86

\* i.e. 5 % of the  $H_s$  values measured in 2003 exceeded 1.37m

### Distribution plots

The distribution of wave parameters are shown in the accompanying graphs of:

- Percentage of occurrence of  $H_s$ ,  $T_p$ ,  $T_z$  and Direction for 2007
- Percentage wave height exceedance (all recorded years) – note that the statistics for 2003 were based on measurements from July to December only
- Joint distribution of all parameters for 2007, given both as number of observations and as percentage of occurrence
- Cumulative joint distribution of parameters from start of records (percentage of occurrence only)
- Incidence of storms during 2007 and for all previous years. Storm events are defined using the Peaks-over-Threshold method. The highest  $H_s$  of each storm event is shown.
- Annual time series of  $H_s$  (red line is storm threshold)






### General

The buoy was first deployed on 08 July 2003. The wave directions recorded by the Datawell Directional WaveRider Mk III were found to be contaminated by a significant tidal signature, compounded by the on-board data processing. The buoy received new electronics to fix this problem in February 2004; wave directions measured before April 2004 were excluded from the analysis.




### Acknowledgements

Tidal data were supplied by the British Oceanographic Data Centre as part of the function of the National Tidal and Sea Level Facility, hosted by the Proudman Oceanographic Laboratory and funded by DEFRA and the Natural Environment Research Council.



Folkestone 2007



Folkestone 2007 - Joint distribution



Folkestone 2007 - Joint distribution (% of occurrence)



Folkestone 2003 to 2007 - Joint distribution (% of occurrence)

